

ATCC

Catalogue of Recombinant DNA Materials 2nd edition, 1991

HOSTS • LIBRARIES • VECTORS • CLONES

Quick Index

HOSTS

Animal cell lines p. 1
Microorganisms p. 4

LIBRARIES p. 13

VECTORS p. 16

CLONES

Index p. 70
Animals p. 81
Bacteria p. 100
Fungi/Yeasts p. 106
Plants p. 110
Viruses p. 111
Viroids p. 125
Unknown p. 132

SACCHAROMYCES

Genomic clone maps p. 133

MEDIA

Formulations p. 197

FORMS

p. 199

INDEXES

By ATCC® Number p. 219



General p. 230

"How to use the catalogue"

is inside the front cover.



American
Type Culture
Collection



removed. Gene (Amst.) 41: 337-342, 1986. (Medium 1236 37C) **Shipped:** in freeze-dried *Escherichia coli* JM101.

pBGS131- (plasmid)

37445 B.G. Spratt. Construction: pBR322, Tn903, M13tg131, bacteriophage f1 *ori*. Size(kb): 4.4. Marker(s): Kan^r. Cloning sites: *EcoRI SmaI SstI EcoRV SphI KpnI XbaI HindIII BamHI AccI HincII SalI PstI BglII*. Replicon(s): pMB1, f1. Contains MCS. Kanamycin-resistant analog of pEMBL. The duplicate *HindIII*, *SmaI* and *AccI* sites have been removed. Gene (Amst.) 41: 337-342, 1986. (Medium 1236 37C) **Shipped:** in freeze-dried *Escherichia coli* JM101.

pBLA 11 (plasmid)

39788 Bio-technology General Corporation. Marker(s): Tet^r. Promoter(s): λ PL. Contains the ribosomal binding site of the β -lactamase gene. U.S. Patent No. 4,742,004 dated May 3, 1988. Note: This material is cited in a U.S. and/or other Patent and may not be used to infringe the patent claims. (Medium 1273 37C) **Shipped:** in freeze-dried *Escherichia coli*.

pBLCAT2 (plasmid)

37527 B. Luckow. Construction: pUC18, SV40, cml, herpes simplex virus *tk* promoter. Size(kb): 4.5. Marker(s): Amp^r. Cloning sites: *HindIII SphI (PstI) SalI XbaI BamHI BglII XhoI ClaI SmaI KpnI SstI (EcoRI)*. Replicon(s): pMB1. Promoter(s): HSV *tk*. Contains MCS. Developed to simplify construction of hybrid CAT genes. Nucleic Acids Res. 15: 5490, 1987. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli*.

pBLCAT3 (plasmid)

37528 B. Luckow. Construction: pUC18, SV40, cml. Size(kb): 4.34. Marker(s): Amp^r. Cloning sites: *HindIII SphI PstI SalI XbaI BamHI BglII XhoI ClaI SmaI KpnI SstI (EcoRI)*. Replicon(s): pMB1. Contains MCS. Developed to simplify construction of hybrid CAT genes. Nucleic Acids Res. 15: 5490, 1987. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli*.

pBR313 (plasmid)

37018 H. Boyer. Other names: NRRL B-14220. Size(kb): 9.6. Marker(s): Amp^r, Tet^r. Cloning sites: *EcoRI SmaI HpaI HindIII BamHI SalI*. Replicon(s): pMB1. A general purpose plasmid vector. Gene (Amst.) 2: 75-93, 1977. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBR322 (plasmid)

31344 Genentech, Inc. Size(kb): 4.363. Marker(s): Amp^r, Tet^r. Cloning sites: *EcoRI ClaI HindIII EcoRV BamHI SphI SalI XmaI NruI Aval BglI PvuII Tth1111 NdeI PstI PvuI ScaI AatII*. Replicon(s): pMB1. A general purpose plasmid vector. U.S. Patent No. 4,366,246 dated Dec. 28, 1982; U.S. Patent No. 4,342,832 dated Aug. 3, 1982. Note: This material is cited in a U.S. and/or other Patent and may not be used to infringe the patent claims. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

37017 Preceptrol® culture. H. Boyer. Construction: pBR313. Size(kb): 4.363. Marker(s): Amp^r, Tet^r. Cloning sites: *EcoRI ClaI HindIII EcoRV BamHI SphI SalI XmaIII NruI Aval BglI PvuII Tth1111 NdeI PstI PvuI ScaI AatII*. Replicon(s): pMB1. A general purpose plasmid vector. Gene (Amst.) 2: 95-113, 1977. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBR327 (plasmid)

37516 E.M. Lederberg. Size(kb): 3.3. Marker(s): Amp^r, Tet^r. Cloning sites: *AatII Aval ClaI EcoRI HgiEII HindIII BamHI EcoRV NruI PstI PvuI SalI ScaI SphI XmaIII XmnI*. Replicon(s): pMB1. A general purpose plasmid vector. Derived from pBR322 by deleting sequences between nt 1430 (*Aval*) and 2519. The *bom* or *Mob* site has been deleted. Recomb. DNA Tech. Bull. 2: 1980; Nature (Lond.) 283: 216-218, 1980; Gene (Amst.) 13: 25-35, 1981; *ibid.*, 50: 3-40, 1986. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* GM21.

pBR328 (plasmid)

37517 E.M. Lederberg. Construction: pBR327 *ori*, pBR325 cml. Size(kb): 4.9. Marker(s): Amp^r, Cml^r, Tet^r. Cloning sites: *AatII AsuII Aval ClaI HgiEII HindIII Tth1111 BamHI EcoRI EcoRV BglI BamHI EcoRI EcoRV NcoI NruI PstI PvuI PvuII SalI SphI XmaIII*. Replicon(s): pMB1. A general purpose plasmid vector. Contains a 482 bp inverted

duplication. Constructed from the *PstI/BamHI* fragment of pBR327 containing the replication origin and the *PstI/BamHI* fragment of pBR325 containing the *cml* gene. Gene (Amst.) 9: 287-305, 1980; *ibid.*, 50: 3-40, 1986. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBR329 (plasmid)

37264 R.L. Rodriguez. Construction: pBR327, pBR328 837 bp cml. Size(kb): 4.2. Marker(s): Amp^r, Cml^r, Tet^r. Cloning sites: *AatI AsuII Aval ClaI HgiEII HindIII Tth1111 BglI BamHI EcoRI EcoRV NcoI NruI PstI PvuI PvuII SalI SphI XmaIII XmnI*. Replicon(s): pMB1. A general purpose plasmid vector. This plasmid does not carry the inverted duplication region found in pBR328. Gene (Amst.) 17: 79-89, 1982; Rodriguez, R.L.; Tait, R.C. Recombinant DNA techniques. Reading, MA: Addison-Wesley; 1983. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBRH1 (plasmid)

37070 R.L. Rodriguez. Construction: pBR316. Size(kb): 7.7. Marker(s): Amp^r. Cloning sites: *EcoRI*. Replicon(s): pMB1. Promoter cloning plasmid vector using expression of tetracycline resistance for selection. Constructed by inserting an oligonucleotide containing an *EcoRI* site into the *HindIII* site of pBR316, inactivating the *tet* promoter. Rodriguez, R.L.; Denhardt, D.T., eds. Vectors: A survey of molecular cloning vectors and their uses. Boston: Butterworth; 1988:pp. 153-177; Gene (Amst.) 7: 271-288, 1979. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBRH3B (plasmid)

37072 R.L. Rodriguez. Construction: pBR316. Size(kb): 7.7. Marker(s): Amp^r. Cloning sites: *EcoRI*. Replicon(s): pMB1. Promoter cloning plasmid vector using expression of tetracycline resistance for selection. Constructed by inserting an oligonucleotide containing an *EcoRI* site into the *HindIII* site of pBR316, inactivating the *tet* promoter. Resistant to low levels of tetracycline without an insert and can be used to clone only strong promoters. Rodriguez, R.L.; Denhardt, D.T., eds. Vectors: A survey of molecular cloning vectors and their uses. Boston: Butterworth; 1988:pp. 153-177; Gene (Amst.) 7: 271-288, 1979. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBRH4 (plasmid)

37071 R.L. Rodriguez. Construction: pBR322. Size(kb): 4. Marker(s): Amp^r. Cloning sites: *EcoRI*. Replicon(s): pMB1. Promoter cloning plasmid vector using expression of tetracycline resistance for selection. Gene (Amst.) 7: 271-288, 1979. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pBRM (plasmid)

37283 J.P. White. Construction: pBR322. Size(kb): 2. Marker(s): Amp^r. Cloning sites: *BamHI EcoRI*. Replicon(s): pMB1. Proc. Natl. Acad. Sci. USA 79: 233-237, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli*.

pBRN3 (plasmid)

37137 D.B. Wilson. Construction: pBR322. Size(kb): 2. Marker(s): Amp^r. Cloning sites: *HindIII EcoRI BglI PstI*. Replicon(s): pMB1. A general purpose plasmid vector useful for construction of shuttle vectors for animal cells. Deletion includes poison sequences near the origin of replication. A pBR322 deletion from near bp 50 to near bp 2400. Plasmid 7: 287-289, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 SK2284.

pBRN/B (plasmid)

37348 M.M. Haltiner. Construction: pBR322. Size(kb): 3.3. Marker(s): Amp^r. Cloning sites: *BglII NruI PstI EcoRI SalI BamHI ClaI EcoRV HindIII NdeI PvuI SnaI SphI Tth1111 XmaI*. Replicon(s): pMB1. Plasmid vector for use in generating nested sets of deletion mutations. Contains a linker cassette sequence containing *NruI* and *BglII* sites adjacent to each other. Nucleic Acids Res. 13: 1015-1025, 1985. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* HB101.

pBRN/S (plasmid)

37347 M.M. Haltiner. Construction: pBR322. Size(kb): 3.3. Marker(s): Amp^r. Cloning sites: *SacI NruI PstI EcoRI SalI BamHI ClaI EcoRV HindIII NdeI PvuI SnaI SphI Tth1111*

pCM4 (plasmid)

37174 R.L. Rodriguez. Construction: pBR327, Tn9 *cml*. Size(kb): 4.2. Marker(s): Amp^r. Replicon(s): pMB1. Contains promoterless CAT gene flanked by *Bam*HI sites which can be used to construct other vectors. Gene (Amst.) 20: 305-316, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pCM7 (plasmid)

37173 R.L. Rodriguez. Construction: pBR327, Tn9 *cml*. Size(kb): 4.2. Marker(s): Amp^r. Replicon(s): pMB1. Contains promoterless CAT gene flanked by *Hind*III sites which can be used to construct other vectors. Gene (Amst.) 20: 305-316, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pcos1EMBL (cosmid)

37568 A. Craig. Construction: R6K, pcos4EMBL. Size(kb): 6.3. Marker(s): Kan^r, Tet^r. Cloning sites: *Bam*HI. Replicon(s): R6K. One of a family of cosmid vectors (ATCC 37568-37571) containing a R6K origin. Contains one λ *cos* site. Gene (Amst.) 57: 229-237, 1987. (Medium 1273 37C) **Shipped:** in freeze-dried *Escherichia coli* DH1.

pcos2EMBL (cosmid)

37569 A. Craig. Construction: pcos1EMBL. Size(kb): 6.1. Marker(s): Kan^r, Tet^r. Cloning sites: *Bam*HI. Replicon(s): R6K. One of a family of cosmid vectors (ATCC 37568-37571) containing a R6K origin. Differs from pcos1EMBL (ATCC 37568) in having 2 λ *cos* sites and a deletion in the region of the Kan^r gene. Gene (Amst.) 57: 229-237, 1987. (Medium 1273 37C) **Shipped:** in freeze-dried *Escherichia coli* DH1.

pcos5EMBL (cosmid)

37570 A. Craig. Construction: pcos3EMBL. Size(kb): 6. Marker(s): Kan^r. Cloning sites: *Bam*HI. Replicon(s): R6K. Contains MCS. One of a family of cosmid vectors (ATCC 37568-37571) containing a R6K origin. Differs from pcos3EMBL in having *Not*I sites flanking the *Bam*HI cloning site. Gene (Amst.) 57: 229-237, 1987. (Medium 1236 37C) **Shipped:** in freeze-dried *Escherichia coli* DH1.

pcos6EMBL (cosmid)

37571 A. Craig. Construction: pcos5EMBL, Pi. Size(kb): 6. Marker(s): Kan^r. Cloning sites: *Bam*HI. Replicon(s): R6K. Contains MCS. One of a family of cosmid vectors (ATCC 37568-37571) containing a mutated R6K origin for increased copy number. Has increased copy number over pcos5EMBL (ATCC 37570). Gene (Amst.) 57: 229-237, 1987. (Medium 1236 37C) **Shipped:** in freeze-dried *Escherichia coli* DH1.

pCR1 (plasmid)

37135 D.R. Helinski. Construction: ColE1. Size(kb): 11.4. Marker(s): Kan^r. Cloning sites: *Eco*RI *Hind*III. Replicon(s): ColE1. A general purpose plasmid vector. Science (Washington, DC) 196: 172-174, 1977. (Medium 1236 37C) **Shipped:** in freeze-dried *Escherichia coli* C600.

pCS3 (plasmid)

39142 Cetus Corporation. Construction: pEW27, pOP9. Marker(s): Amp^r, Tet^r. Cloning sites: *Eco*RI *Pvu*II *Pst*I *Pvu*I *Clal* *Hind*III *Sph*I *Sac*I *Bam*HI *Nru*I. Replicon(s): ColE1. Gives high copy number at high temperature. Contains the replication origin for high copy number at high temperature of pEW27 (ATCC 37124). pOP9 was constructed by cloning the *Eco*RI/*Pvu*II fragment containing the origin of pOP6 into pBR322. This should NOT be grown over 30C; it is best between 28-30C. U.S. Patent No. 4,677,064 dated June 30, 1987; Proc. Natl. Acad. Sci. USA 79: 3570-3574, 1982. Note: This material is cited in a U.S. and/or other Patent and may not be used to infringe the patent claims. (Medium 1227 28C) **Shipped:** in freeze-dried *Escherichia coli* MM294.

pD553 (plasmid)

37284 D.J. Drahos. Construction: pKO3, λ . Marker(s): Amp^r. Cloning sites: *Eco*RI *Hind*III *Hpa*I. Replicon(s): pMB1. Terminator: λ *tL1*. Plasmid containing antiterminating λ gene N and *nut L* site. Gene (Amst.) 16: 261-274, 1981. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* C600 *galK*⁻.

pDG141 (plasmid)

39588 Cetus Corporation. Other names: CMCC 1966. Construction: pBR322. Size(kb): 2.5. Marker(s): Amp^r. Cloning sites: *Sac*I. Replicon(s): pMB1. Promoter(s): *trp*.

Expression vector with a tryptophan promoter-operator and ribosome binding site operably linked with an ATG start codon. U.S. Patent No. 4,889,818 dated Dec. 26, 1989; U.S. Patent No. 4,711,845 dated Dec. 12, 1987; U.S. Patent No. 4,784,949 dated Nov. 15, 1988. Note: This material is cited in a U.S. and/or other Patent and may not be used to infringe the patent claims. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli*.

pDM1 (plasmid)

37357 D. Mead. Construction: bacteriophage f1, pAT153. Size(kb): 3.8. Marker(s): Amp^r. Replicon(s): pMB1. Contains an easily isolated f1 intergenic region for conversion of any plasmid to an ssDNA vector (use *Sac*I plus *Acc*I plus *Hinc*II). Nucleic Acids Res. 13: 1103-1118, 1985. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* NM522.

pDR42 (plasmid)

37180 G.N. Bennett. Construction: pBR322, *Escherichia coli trp*. Size(kb): 4.4. Marker(s): Amp^r, Tet^r. Cloning sites: *Pst*I *Eco*RI *Hind*III *Bam*HI *Sac*I. Replicon(s): pMB1. A pBR322 derivative with increased tetracycline resistance and increased fusaric acid sensitivity. Hybrid *trp-tet* promoter formed by insertion of partial *trp* promoter at the *Clal* site. Plasmid 7: 290-293, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* RR1.

pDR540 (plasmid)

37282 G.N. Bennett. Construction: pKO1, *Escherichia coli trp-lac* fusion promoter *tac*. Marker(s): Amp^r, *galK*⁺. Replicon(s): pMB1. Promoter(s): *tac*. Contains an easily purifiable *trp-lac* hybrid promoter which can be used to construct expression vectors. Unstable when transformed into C600 *galK*⁻. Gene (Amst.) 20: 231-243, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* JM103.

pEA300 (plasmid)

37181 E. Amann. Construction: pKK84-1, *ptrp*H1. Size(kb): 5.5. Marker(s): Amp^r. Cloning sites: *Eco*RI *Clal* *Hind*III *Bam*HI *Pvu*II. Replicon(s): pMB1. Promoter(s): *trp*. Terminator: *rrnB*. For construction of -35 *trp* expression vectors. Constructed by inserting a 192 bp fragment containing the -35 sequence of the *trp* promoter into the *Clal* site of pKK84-1. Contains two terminators tandemly arranged. Maniatis, T. et al., eds. Molecular cloning: a laboratory manual. Cold Spring Harbor, NY: CSHL; 1982:pp. 413-446; Gene (Amst.) 25: 167-178, 1983. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* W3110 *lacI*^q.

pEMBL8⁺ (phagemid)

37396 G. Cesareni. Construction: pUC8, bacteriophage f1. Size(kb): 3.99. Marker(s): Amp^r. Cloning sites: *Eco*RI *Ava*I *Sma*I *Xma*I *Bam*HI *Sac*I *Acc*I *Pst*I *Hind*II *Hind*III *Clal* *Eco*B *Nae*I. Replicon(s): pMB1, f1. Contains MCS. ssDNA-producing plasmid with polylinker in *lacZ'*. Nucleic Acids Res. 11: 1645-1655, 1983. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 71/18.

pEMBL8⁻ (plasmid)

37397 G. Cesareni. Construction: pUC8, bacteriophage f1. Size(kb): 3.99. Marker(s): Amp^r. Cloning sites: *Eco*RI *Ava*I *Sma*I *Xma*I *Bam*HI *Sac*I *Acc*I *Pst*I *Hind*II *Hind*III *Clal* *Eco*B *Nae*I. Replicon(s): pMB1, f1. Contains MCS. ssDNA-producing plasmid with polylinker in *lacZ'*. Nucleic Acids Res. 11: 1645-1655, 1983. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 71/18.

pEMBL9 (plasmid)

37395 G. Cesareni. Construction: pUC9, pEMBL8, *lacZ*. Size(kb): 3.99. Marker(s): Amp^r. Cloning sites: *Eco*RI *Sma*I *Bam*HI *Sac*I *Ava*I *Pst*I *Hind*III. Contains MCS. ssDNA-producing plasmid with polylinker in *lacZ'*. Nucleic Acids Res. 11: 1645-1655, 1983; Gene (Amst.) 35: 27-32, 1985. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 71/18.

pEMBL18-Not (Sma-) (plasmid)

37573 A. Craig. Construction: pEMBL18. Size(kb): 3.9. Marker(s): Amp^r. Cloning sites: *Eco*RI *Sac*I *Kpn*I *Not*I *Bam*HI *Sac*I *Pst*I *Sph*I *Hind*III. Replicon(s): pMB1. Promoter(s): *lac*. Contains MCS. General purpose vector modified to include a *Not*I site to facilitate library construction and chromosome walking. *Sma*I site in polylinker of pEMBL18 changed to a *Not*I site. Gene (Amst.) 57: 229-237, 1987. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* JM101.

pLA7 (plasmid)

37281 I.R. Beacham. Construction: pCB1, pBR322. Size(kb): 4.1. Marker(s): Amp^r, *ush*. Cloning sites: *Bcl*I *Ava*I. Replicon(s): pMB1. A positive selection vector for cloning *Sau*3AI-generated DNA fragments. Growth of the plasmid in a *dam*⁻ host is necessary for *Bcl*I cleavage. Gene (Amst.) 27: 323-325, 1984. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* GM33.

pLG200 (plasmid)

37113 L.P. Guarente. Construction: pBR322, *Escherichia coli*. Size(kb): 8.9. Marker(s): Amp^r. Cloning sites: *Hind*III. Replicon(s): pMB1. Promoter(s): *lacUV5*. Can be used to construct a plasmid which directs the expression of a cloned gene under the control of the *lacUV5* promoter. Cell 20: 543-553, 1980. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* LG90.

pLG338 (plasmid)

37130 N.G. Stoker. Construction: pSC105. Size(kb): 7.3. Marker(s): Kan^r, Tet^r. Cloning sites: *Kpn*I *Eco*RI *Xho*I *Sma*I *Bam*HI *Hinc*II/*Sal*I. Replicon(s): pSC101. A low copy number (6-8 per chromosome), general purpose plasmid vector. Gene (Amst.) 18: 335-341, 1982. (Medium 1273 37C) **Shipped:** in freeze-dried *Escherichia coli* C600.

pLG339 (plasmid)

37131 N.G. Stoker. Construction: pSC105. Size(kb): 6.3. Marker(s): Kan^r, Tet^r. Cloning sites: *Pvu*II *Eco*RI *Xho*I *Sma*I *Bam*HI *Sph*I *Hinc*II/*Sal*I. Replicon(s): pSC101. A low copy number (6-8 per chromosome), general purpose plasmid vector. Gene (Amst.) 18: 335-341, 1982. (Medium 1273 37C) **Shipped:** in freeze-dried *Escherichia coli* C600.

pLG400 (plasmid)

37114 L.P. Guarente. Construction: pBR322, *Escherichia coli*. Size(kb): 8.9. Marker(s): Amp^r. Cloning sites: *Hind*III *Bam*HI. Replicon(s): pMB1. Promoter(s): *lacUV5*. Can be used to construct a plasmid which directs the expression of a cloned gene under the control of the *lacUV5* promoter. Cell 20: 543-553, 1980. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* LG90.

pLKC480 (plasmid)

37594 A.A. Tiedman. Construction: A *lacZY* plasmid, Tn5. Size(kb): 9.9. Marker(s): Amp^r, Kan^r. Cloning sites: *Eco*RI *Bam*HI *Sal*I *Hind*III. Replicon(s): pMB1. Contains MCS. One of a series of *lacZY* fusion vectors (ATCC 37594-37596) that allow fusions in one of three reading frames and retain flanking DNA for homologous recombination into the *Escherichia coli* genome. The 6.3 kb *lacZY*-Kan^r cassette is released with *Sma*I. A 1.3 kb *Hind*III/*Nru*I Kan^r fragment (with *Hind*III site filled in) was inserted at the *Nru*I site of a *lacZY* fusion plasmid. Nucleic Acids Res. 16: 3587, 1988. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* TX528.

pLKC481 (plasmid)

37595 A.A. Tiedman. Construction: A *lacZY* plasmid, Tn5. Size(kb): 9.9. Marker(s): Amp^r, Kan^r. Cloning sites: *Eco*RI *Bam*HI *Sal*I *Hind*III. Replicon(s): pMB1. Contains MCS. One of a series of *lacZY* fusion vectors (ATCC 37594-37596) that allow fusions in one of three reading frames and contain flanking DNA for homologous recombination into the *Escherichia coli* genome. The 6.3 kb *lacZY*-Kan^r cassette is released with *Sma*I. A 1.3 kb *Hind*III/*Nru*I Kan^r fragment (with *Hind*III site filled in) was inserted at the *Nru*I site of a *lacZY* fusion plasmid. Nucleic Acids Res. 16: 3587, 1988. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli*.

pLKC482 (plasmid)

37596 A.A. Tiedman. Construction: A *lacZY* plasmid, Tn5. Size(kb): 9.9. Marker(s): Amp^r, Kan^r. Cloning sites: *Eco*RI *Bam*HI *Sal*I *Hind*III. Replicon(s): pMB1. Contains MCS. One of a series of *lacZY* fusion vectors (ATCC 37594-37596) that allow fusions in one of three reading frames and retain flanking DNA for homologous recombination into the *Escherichia coli* genome. The 6.3 kb *lacZY*-Kan^r cassette is released with *Sma*I. A 1.3 kb *Hind*III/*Nru*I Kan^r fragment (with *Hind*III site filled in) was inserted at the *Nru*I site of a *lacZY* fusion plasmid. Nucleic Acids Res. 16: 3587, 1988. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* TX528.

pLM0.8 (plasmid)

39604 Actagen, Inc. Marker(s): Amp^r. Note: This material is cited in a U.S. and/or other Patent and may not be used to infringe the patent claims. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* HB101.

pLV57 (plasmid)

37177 D. O'Connor. Construction: pBR325, NTP14m07(Ts). Size(kb): 6.1. Marker(s): Amp^r, Cml^r, *Eco*RI m(Ts)^r. Cloning sites: *Hind*III *Bgl*II *Ava*I *Eco*RI. Replicon(s): pMB1. Positive selection cloning vector employing *Eco*RI restriction-modification system. Gene (Amst.) 20: 219-229, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* HB101.

pLV59 (plasmid)

37178 D. O'Connor. Construction: pACYC184, NTP14m(Ts). Size(kb): 6.3. Marker(s): Cml^r, *Eco*RI m(Ts)^r, Tet^r. Cloning sites: *Hind*III *Bgl*II *Pst*I *Bam*HI *Eco*RI *Sal*I/*Hinc*II *Hinc*II. Replicon(s): p15A. Positive selection cloning vector employing *Eco*RI restriction system. Gene (Amst.) 20: 219-229, 1982. (Medium 1675 37C) **Shipped:** in freeze-dried *Escherichia coli* HB101.

pMAM17 (plasmid)

37325 B. Polisky. Construction: pPLc236, ColE1 *rop*. Size(kb): 3.01. Marker(s): Amp^r. Cloning sites: *Eco*RI *Bam*HI *Hind*III *Pvu*II. Replicon(s): ColE1. Promoter(s): λ PL. Expression vector that permits positive selection of DNA inserts; also permits high level temperature-induced expression of inserted DNA as fusion protein. Gene (Amst.) 31: 155-164, 1984. (Medium 1227 30C) **Shipped:** in freeze-dried *Escherichia coli* K-12 Δ H1 Δ trp.

pMB9 (plasmid)

37019 H. Boyer \leftarrow F. Bolivar. Construction: pMB8, pSC101. Size(kb): 5.2. Marker(s): Tet^r. Cloning sites: *Bam*HI *Hind*III *Sal*I *Eco*RI. Replicon(s): pMB1. A general purpose plasmid vector. Gene (Amst.) 2: 75-93, 1977. (Medium 1273 37C) **Shipped:** in freeze-dried *Escherichia coli* HB101.

pMF7 (cosmid)

37117 M. Feiss. Construction: pBR322, λ . Size(kb): 6.8. Marker(s): Amp^r. Cloning sites: *Eco*RI *Sal*I. Replicon(s): pMB1. A general purpose cosmid vector. Gene (Amst.) 17: 123-130, 1982; Proc. Natl. Acad. Sci. USA 79: 3498-3502, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 IC202.

pMF517 (cosmid)

37118 M. Feiss. Construction: pBR322, λ . Size(kb): 7.1. Marker(s): Amp^r. Cloning sites: *Clal* *Hind*III *Sal*I *Pst*I *Eco*RI. Replicon(s): pMB1. A general purpose cosmid vector. Gene (Amst.) 17: 123-130, 1982; Proc. Natl. Acad. Sci. USA 79: 3498-3502, 1982. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 IC202.

pMH621 (plasmid)

31775 NCI-Frederick Cancer Research Center. Size(kb): 7.6. Marker(s): Amp^r. Cloning sites: *Bgl*II. Replicon(s): pMB1. Promoter(s): *ompF*. An expression vector for cytoplasmic export of a cloned gene product. Maniatis, T.; *et al.*, eds. Molecular cloning: A laboratory manual. Cold Spring Harbor, NY: CSHL; 1982:pp. 429-430. Note: This material is cited in a U.S. and/or other Patent Application and may not be used to infringe the patent claims. (Medium 1227 37C) **Shipped:** in freeze-dried *Escherichia coli* K-12 MH2000.

40038 NCI-Frederick Cancer Research Center. Size(kb): 7.6. Marker(s): Amp^r. Cloning sites: *Bgl*II. Replicon(s): pMB1. Promoter(s): *ompF*. An expression vector for cytoplasmic export of cloned gene product by formation of a fusion protein with the *ompF* signal sequence. Requires as host MH3000 (ATCC 35468) for viability and TK 1046 (ATCC 35467) for protein expression. Note: This material is cited in a U.S. and/or other Patent Application and may not be used to infringe the patent claims. **Shipped:** as thawed purified plasmid DNA, supplied by depositor.

pMLB1034 (plasmid)

37222 M.L. Berman. Construction: pMC871, pBR322. Size(kb): 7.2. Marker(s): Amp^r. Cloning sites: *Bam*HI *Eco*RI *Sma*I. Replicon(s): pMB1. Contains MCS. For screening promoters by determining β -galactosidase activity (*lacZ*). The cloning sites are 5' to a promoterless β -galactosidase gene. The insert

EUKARYOTIC GENES CLONED AND EXPRESSED PRIOR TO MARCH 19, 1980

	Gene	Reference	System
MAMMALIAN GENE: β -GALACTOSIDASE EXPRESSION SYSTEM			
1	human somatostatin	Itakura et al. (1977)	β -gal promoter
2	chicken ovalbumin	Fraser et al. (1978) Mercereau et al. (1978)	β -gal promoter
3	matute human insulin	Goeddel et al. (1979a)	β -gal promoter
MAMMALIAN GENE: β -LACTAMASE EXPRESSION SYSTEM (Inserted into PstI site of pBR322)			
4	rat preproinsulin	Villa-Komaroff (1978)	<i>bla</i> promoter
5	mouse dihydrofolate reductase	Chang et al. (1978)	<i>bla</i> promoter
6	rat pregrowth hormone	Seeburg et al. (1978)	<i>bla</i> promoter
MAMMALIAN GENE: <i>TRP D</i> EXPRESSION SYSTEM			
7	human pregrowth hormone	Martial et al. (1979)	<i>trp</i> promoter
MAMMALIAN GENE: <i>LAC</i> PROMOTER			
8	human mature growth hormone	Goeddel et al. (1979b)	<i>lac</i> promoter ¹
MAMMALIAN GENE: OTHER			
9	human globin	Wilson et al. (1979)	...
VIRAL GENE (Genes which are expressed normally by infected eukaryotic hosts)			
10	simian virus 40 t antigen	Roberts et al. (1979)	<i>lac</i> promoter ²
11	human hepatitis B virus	Burrell et al. (1979)	<i>bla</i> promoter
12	fowl plague virus	Emtage et al. (1980)	<i>trp</i> promoter Insert at pBR322 HindIII
YEAST GENE			
13	<i>Neurospora</i> dehydroquinolate hydrolase	Vapnek et al. (1977)	Insert at pBR322 HindIII/EcoRI
14	<i>Saccharomyces His and Leu</i>	Ratzkin et al. (1977)	ColEI plasmid
15	<i>Saccharomyces</i> galactokinase	Schell et al. (1979)	Insert at pBR322 BamHI
16	<i>Saccharomyces</i> OMP decarboxylase	Bach et al. (1979)	Insert at pBR322 HindIII

Key: β -gal = β -galactosidase gene
bla = β -lactamase gene

trp = tryptophan operon
lac = lactose operon

¹ System subsequently used by Goeddel et al. (Nucleic Acids Research, Vol. 8, No. 18, pp. 4057-4074 (1980)), to express IFN- β .

² System subsequently used by Taniguchi et al. (PNAS, 77:5230-5233 (September 1980)), to express IFN- β .

LIST OF CITES

1. Itakura et al., Science, 198:1056-1063 (December 1977)
2. Fraser et al., PNAS, 75:5936-5940 (December 1978)
Mercereau-Puijalon et al., Nature, 275:505-510 (October 12, 1978)
3. Goeddel et al., PNAS, 76:106-110 (January 1979a)
4. Villa-Komaroff et al., PNAS 75:3727-3731 (August 1978)
5. Chang et al., Nature, 275:617-624 (October 19, 1978)
6. Seeburg et al., Nature, 276:795-798 (December 21/28, 1978)
7. Martial et al., Science, 205:602-607 (August 1979)
8. Goeddel et al., Nature, 281:544-548 (October 18, 1979b)
9. Wilson et al., PNAS, 76:5631-5635 (1979)
10. Roberts et al., PNAS, 76:5596-5600 (November 1979)
11. Burrell et al., Nature, 279:43-47 (May 3, 1979)
12. Emtage et al., Nature, 283:171-174 (January 10, 1980)
13. Vapnek et al., PNAS, 74:3508-3512 (August 1977)
14. Ratzkin et al., PNAS, 74:487-491 (February 1977)
15. Schell et al., Gene, 5:291-303 (1979)
16. Bach et al., PNAS, 76:386-390 (January 1979)